



SPECTROPHOTOMETRIC ANALYSIS OF FULVIC ACID SOLUTIONS – A SECOND LOOK

Elham A. Ghabbour* and Geoffrey Davies

Department of Chemistry & Chemical Biology, Northeastern University, Boston, MA 02115, USA

Received April 7, 2009; in final form June 8, 2009; Accepted June 11, 2009.

SUPPLEMENTARY MATERIALS

Figure S1 Plots of Eq. (2) for IHSS standard fulvic acids **1F**, **2F** and **3F** at (a) pH 1.0 and (b) pH 4.0.

Figure S2 Plots of absorbance vs. FA concentration for IHSS standard fulvic acids **1F**, **2F** and **3F** at (a) pH 1.0 and (b) pH 4.0. The slope of each plot is the absorption coefficient ϵ .

* Corresponding author: Department of Chemistry & Chemical Biology, Northeastern University, Boston, MA 02115, USA. Phone: 617-373-7988; fax: 617-373-8795, e-mail: <e.ghabbour@neu.edu>.




Figure S1 Plots of Eq. (2) for IHSS standard fulvic acids 1F, 2F and 3F at pH 1.0 and pH 4.0.




Figure S2 Plots of absorbance vs. FA concentration for IHSS standard fulvic acids 1F, 2F and 3F at (a) pH 1.0 and (b) pH 4.0. The slope of each plot is the absorption coefficient ϵ .




Figure S2 Plots of absorbance vs. FA concentration for IHSS standard fulvic acids 1F, 2F and 3F at pH 1.0 and pH 4.0. The slope of each plot is the absorption coefficient ϵ .

AES 9407

© Northeastern University, 2009